

Trial record **1 of 1** for: Vargatef in 2nd-line Therapy of Non-Small Cell Lung Cancer (NSCLC)

[Previous Study](#) | [Return to List](#) | [Next Study](#)

VARGADO - Vargatef in 2nd-line Therapy of Non-Small Cell Lung Cancer (NSCLC)

This study is currently recruiting participants. (see [Contacts and Locations](#))

Verified November 2016 by Boehringer Ingelheim

Sponsor:

Boehringer Ingelheim

Information provided by (Responsible Party):

Boehringer Ingelheim

ClinicalTrials.gov Identifier:

NCT02392455

First received: March 17, 2015

Last updated: November 21, 2016

Last verified: November 2016

[History of Changes](#)

[Full Text View](#)

[Tabular View](#)

[No Study Results Posted](#)

[Disclaimer](#)

[How to Read a Study Record](#)

Purpose

This observational study will investigate the efficacy and tolerability of **Vargatef (Nintedanib)** plus docetaxel in daily routine second-line **treatment** in patients with locally advanced, metastatic or locally recurrent **NSCLC**. **Treatment with Vargatef** in eligible **NSCLC** patients, for whom the treating physician has decided to initiate **treatment with Vargatef** in second line according to the local label, will be observed for up to 24 months. Survival follow-up will be done until the end of the study.

Condition	Intervention
Carcinoma, Non-Small-Cell Lung	Drug: treatment

Study Type: Observational

Study Design: Observational Model: Cohort

Time Perspective: Prospective

Official Title: **Vargatef in 2nd-line Therapy** of Advanced or Metastatic Adenocarcinoma of the **Lung**

Resource links provided by NLM:

[Genetics Home Reference](#) related topics: [lung cancer](#)

[Drug Information](#) available for: [Nintedanib](#) [Nintedanib esylate](#)

[Genetic and Rare Diseases Information Center](#) resources: [Lung Adenocarcinoma](#)

[U.S. FDA Resources](#)

Further study details as provided by Boehringer Ingelheim:

Primary Outcome Measures:

- Percentage of patients alive one year after start of **therapy with Vargatef** and Docetaxel (1-year survival rate) [Time Frame: up to 24 months]
[Designated as safety issue: No]

Secondary Outcome Measures:

- One year survival rate of patients with first line progression within 9 months after start of first line **therapy** [Time Frame: up to 24 months]
[Designated as safety issue: No]
- Progression-free survival of patients with first line progression within 9 months after start of first line **therapy** [Time Frame: up to 24 months]
[Designated as safety issue: No]
- Median overall survival [Time Frame: up to 24 months] [Designated as safety issue: No]
- Progression-free survival [Time Frame: up to 24 months] [Designated as safety issue: No]
- Tumour** control rate (complete response, partial response, stable disease) [Time Frame: up to 24 months] [Designated as safety issue: No]
- Incidence of side effects [Time Frame: up to 24 months] [Designated as safety issue: Yes]

Estimated Enrollment: 400
Study Start Date: March 2015
Estimated Study Completion Date: December 2020
Estimated Primary Completion Date: March 2019 (Final data collection date for primary outcome measure)

<u>Groups/Cohorts</u>	<u>Assigned Interventions</u>
A	Drug: treatment Docetaxel plus nintedanib until progression or intolerance

▶ Eligibility

Ages Eligible for Study: 18 Years and older (Adult, Senior)
Genders Eligible for Study: Both
Accepts Healthy Volunteers: No
Sampling Method: Probability Sample

Study Population

NSCLC patients

Criteria

Inclusion criteria:

- age 18 or older
- men and women locally advanced, metastatic and/or recurrent NSCLC with adenocarcinoma histology for which vargatef treatment is indicated according to Summary of Product Characteristics (SmPC)
- after first line therapy; second-line therapy not yet started
- standard 21-day-cycles docetaxel treatment according to SmPC possible
- written informed consent

Exclusion criteria:

- contraindications according to the SmPC of Vargatef or Docetaxel
- more than one chemotherapy for treatment of NSCLC in palliative setting
- current participation in a clinical trial
- pregnancy
- breastfeeding

▶ Contacts and Locations

Choosing to participate in a study is an important personal decision. Talk with your doctor and family members or friends about deciding to join a study. To learn more about this study, you or your doctor may contact the study research staff using the Contacts provided below. For general information, see [Learn About Clinical Studies](#).

Please refer to this study by its ClinicalTrials.gov identifier: NCT02392455

Contacts

Contact: Boehringer Ingelheim Call Center 1-800-243-0127 clintrriage.rdg@boehringer-ingelheim.com

Locations

Germany

Boehringer Ingelheim Investigational Site **Recruiting**
one or multiple Investigational sites, Germany

Sponsors and Collaborators

Boehringer Ingelheim

Investigators

Study Chair: [Boehringer Ingelheim](#) [Boehringer Ingelheim](#)

▶ More Information

Responsible Party: [Boehringer Ingelheim](#)
ClinicalTrials.gov Identifier: [NCT02392455](#) [History of Changes](#)
Other Study ID Numbers: 1199.211
Study First Received: March 17, 2015

Last Updated: November 21, 2016
Health Authority: Germany: Ethics Commission

Additional relevant MeSH terms:

Carcinoma, **Non-Small-Cell Lung**

Bronchial **Neoplasms**

Lung Neoplasms

Respiratory Tract **Neoplasms**

Thoracic **Neoplasms**

Neoplasms by Site

Neoplasms

Lung Diseases

Nintedanib

Carcinoma, Bronchogenic

Respiratory Tract Diseases

Antineoplastic Agents

Enzyme Inhibitors

Molecular Mechanisms of Pharmacological Action

ClinicalTrials.gov processed this record on November 22, 2016